

The Boyce Clinic Protocols for using Oxygen Therapies to successfully treat HIV+ AIDS patients.

Dr James O. Boyce, of The Boyce Clinic, in Diamondhead, Mississippi USA, has been using oxidative therapies to successfully treat HIV+ and AIDS patients.

He has come under pressure from the medical bureaucracy in the USA, to cease operations and to not disclose his protocols.

Nexus Magazine is pleased to reproduce, with permission, the most recent protocols used by Dr Boyce to successfully treat these diseases. (Also see 'Letters to the Editor')

BOYCE CLINIC, INC. OXIDATIVE PROTOCOL FOR HIV-POSITIVE/AIDS PATIENTS

PRE-ENTRANCE:

- Physical examination
- Blood chemistry examination
- Examination of patient history in detail
- Past and present social history
- Brief psychological questionnaire
- Signed agreement that protocol will be followed while under treatment

DIET:

- No meat, chicken or shellfish.
- Eat raw or slightly steamed vegetables, and natural fruits such as oranges, pears, apples, carrots, etc.
- Juice from fresh, live, whole plants such as celery, carrots, cantaloupe (wash outside with mild, dilute hydrogen peroxide solution to remove possible chemical sprays), etc.
- One regular (junk) meal every week.
- Absolutely no alcohol or drugs to be taken (other than prescribed) as they will

negate the therapy. Note: One beer shuts down the immune system for a minimum of 24 hours.

TREATMENTS

Minor Autohemotherapy:

Take 3 cc of patient's blood, mix with 21 cc of 210 micrograms of ozone and re-inject into the vein.

IV Peroxide (white bag):

To a 250 cc bag of 5% dextrose (use ringer solution if patient is diabetic) add 5 cc of sodium bicarbonate, 5 cc of magnesium chloride, 2 cc of 0.375% solution of hydrogen peroxide, 2 cc of 99% DMSO and 5 cc of lidocaine.

This IV drip should be given over a period of 1 1/2 to 2 hours. If patient experiences discomfort then check the IV needle for position.

Direct IV Ozone:

After 3 days of minor autohemotherapy, then in addition to this procedure, commence with direct IV Ozone treatment by adding 30 cc of 210 MCG ozone, and increasing the volume to approximately 180 cc over the next 7 days. Note: Gradually increase the dosage from 30 cc to 180 cc of medically derived ozone according to patient's comfort levels. Watch for tightness or coughing in chest and/or throat area--indicating blood system is full. When tightness or coughing appears, immediately discontinue ozone injections. Ozone may be injected twice a day at 8 hour intervals.

IV Chelation (pink bag):

The white bag mixture (peroxide) is given 3 days in a row, then on the 4th day, the IV chelation is administered to remove excess free radicals which are created by the infusion of peroxide and ozone. In preparation of the pink bag, the following procedure is followed: 80 cc of solution is removed from a 500 cc bag of 5% dextrose or ringers. Then, 3 grams of EDTA is added along with 50 cc of injectable vitamin C (25 grams per bottle), 5 cc of magnesium chloride, 5 cc of

sodium bicarbonate, 5 cc of B12, 1 cc of B6 (pyroxidine), 1 cc of B1 (Thiamine), and 1 cc of multiple trace elements. Since chelation reduces the blood-sugar level and may cause venule cramping, then up to 10 cc of lidocaine may be added to relieve this discomfort. If a patient notices a drop of sugar level (i.e., lack of energy), then he may eat a piece of candy to raise the blood-sugar level. The pink IV drip should last at least 4 hours, 5 hours being preferred.

Bath Therapy:

First week, the patient should fill the bathtub full of water and add one gallon of 35% hydrogen peroxide. The 2nd week and beyond, the patient should use one quart of 35% hydrogen peroxide and in both cases should soak up to 30 minutes. The patient must be aware not to get the bath water in his eyes, nose, mouth, or ears. Shower afterwards. The skin is the largest organ of the body and it excretes as well as absorbs, therefore the hydrogen peroxide helps to detoxify any toxins that are excreted by the skin.

Colonics:

To assist detoxification of rapid-oxidising viruses, colonics should be given 3 days in a row each week, then once a week for eight weeks, then at least once a month.

Acupuncture:

Needle or laser acupuncture should be used 3 times a week on points G36 and S37

Rest:

Patient should get plenty of rest: no late-night partying with others. Patient should sleep under a pinkish-purple coloured light (non-painted, clear bulb).

Supplementation Therapy:

Twenty-five drops stabilised oxygen in an 8 ounce glass of water or orange juice twice a day, morning and night. There are many brands of stabilised oxygen on the market, such as Aerobic 07 (TM). A less expensive approach would be the utilisation of one drop of hydrogen peroxide to a glass of water or orange juice, morning and night, increasing it by one drop each day. If nausea appears, back off 3 drops and then continue to a maximum of 25 drops, twice a day. By using hydrogen peroxide, you should take acidophilus capsules or powder two hours beforehand.

Genetically engineered wheat grass, source of super-oxide dismutases, tablets

with catalase and/or coenzyme Q10 should be taken twice a day. Example: 6 tablets morning, 6 tablets night, of AOX/Plax by Biomed Foods. Take 5 one-gram tablets of vitamin C twice a day. Also take a multi-vitamin with mineral and an additional B complex tablet once a day. Take a homeopathic Thymus under the tongue, one eye dropper 3 times a day. Example: N-15 Thymus Plus distributed by Progressive Laboratories, Inc., Irving Texas.

Optional Therapy:

It has also been suggested that the patient be put in a hyperbaric oxygen chamber for 1 hour, twice daily, 3 times a week, at 2 ATA. There is also some medical evidence indicating positive results for patients sleeping with the north pole of a 5000 gauss magnet beneath the coccyx (Charlie Farr, MD).

Further studies have suggested (though not yet confirmed) that Aloe Vera, Reshi mushrooms, shark cartilage and Carnivora will raise the T4 count. There is evidence that the Rief/Curtis equipment will kill the "AIDS Virus" when set at 32,750, 33,750, and 34,750 MHz for a period of 10 minutes at each setting. It has been reported that the virus replicates at 31,750 MHz. When using this equipment, colonic irrigation is absolutely necessary to maintain a clean colon.

BOYCE CLINIC, INC.

4410-B West Aloha Dr.,
Diamondhead, MS 39525
USA

Ph: (601) 255 9457
Fx: (601) 255 6591

HYDROGEN PEROXIDE

35% Food Grade Hydrogen Peroxide must be a) handled carefully (direct contact will burn the skin - immediate flushing with water is recommended); b) diluted properly before use; c) stored safely and properly (after making a dilution the remainder should be stored tightly sealed in the freezer).

One of the most convenient methods of dispensing 35% H₂O₂ is from a small glass eye dropper bottle. Fill this with the 35% H₂O₂ and store the larger container in the freezer until more is needed. Store the eye dropper bottle in the refrigerator.

The generally recommended dosage is outlined in the chart below. The drops are mixed with either 6 to 8 ounces of distilled water, juice, milk or even aloe vera juice or gel. (Don't use chlorinated tap water to dilute the peroxide!)

The program outlined here is only a suggestion, but it is based on years of experience, and reports from thousands of users. Those who choose to go at a slower pace can expect to progress more slowly, but that certainly is an option. The program is not carved in stone, and keep in mind that it can be adapted to fit individual needs.

Individuals who have had transplants should not undertake an H₂O₂ programme. Hydrogen Peroxide stimulates the immune system, and could possibly cause a rejection of the organ.

Maintenance Dosage

In most situations after the 21 day programme listed below, the amount of H₂O₂ can be tapered off gradually as follows:

- 25 drops once every other day for 1 week
- 25 drops once every third day for 2 weeks
- 25 drops once every fourth day for 3 weeks

This can then be reduced to between 5 and 15 drops per week based on how one feels.

Those with more serious problems will often benefit from staying on 25 drops three times a day for one to three weeks, then tapering down to 25 drops two times daily until the problem is resolved (possibly as long as six months).

It is important that H₂O₂ be taken on an empty stomach. This is best accomplished by taking it either one hour before meals, or three hours after meals. If there is food in the stomach the reaction of the H₂O₂ on any bacteria present may cause excess foaming, indigestion and possibly even vomiting.

It is recommended that you start taking Vitamin E and an acidophilus product before going onto H₂O₂.

Before you commence a course of self treatment using hydrogen peroxide, we strongly recommend that you become more informed on the subject. To this end, we suggest you obtain the Nexus information kit on Oxygen Therapies (see page 39).

Day #	Number of drops	Times Per Day
1	3	3
2	4	3
3	5	3
4	6	3
5	7	3
6	8	3
7	9	3
8	10	3
9	12	3
10	14	3
11	16	3
12	18	3
13	20	3
14	22	3
15	24	3
16-21	25	3